



HEALTH

Vast Study Casts Doubts on Value of Mammograms

By GINA KOLATA FEB. 11, 2014

One of the largest and most meticulous studies of mammography ever done, involving 90,000 women and lasting a quarter-century, has added powerful new doubts about the value of the screening test for women of any age.

It found that the death rates from breast cancer and from all causes were the same in women who got mammograms and those who did not. And the screening had harms: One in five cancers found with mammography and treated was not a threat to the woman's health and did not need treatment such as chemotherapy, surgery or radiation.

The study, published Tuesday in The British Medical Journal, is one of the few rigorous evaluations of mammograms conducted in the modern era of more effective breast cancer treatments. It randomly assigned Canadian women to have regular mammograms and breast exams by trained nurses or to have breast exams alone.

Researchers sought to determine whether there was any advantage to finding breast cancers when they were too small to feel. The answer is no, the researchers report.

The study seems likely to lead to an even deeper polarization between those who believe that regular mammography saves lives, including many breast cancer patients and advocates for them, and a growing number of researchers who say the evidence is lacking or, at the very least, murky.

"It will make women uncomfortable, and they should be uncomfortable," said Dr. Russell P. Harris, a screening expert and professor of medicine at the University of North Carolina, Chapel Hill, who was not involved in the study. "The decision to have a mammogram should not be a slam dunk."

The findings will not lead to any immediate change in guidelines for

mammography, and many advocates and experts will almost certainly dispute the idea that mammograms are on balance useless, or even harmful.

Dr. Richard C. Wender, chief of cancer control for the American Cancer Society, said the society had convened an expert panel that was reviewing all studies on mammography, including the Canadian one, and would issue revised guidelines later this year. He added that combined data from clinical trials of mammography showed it reduces the death rate from breast cancer by at least 15 percent for women in their 40s and by at least 20 percent for older women.

That means that one woman in 1,000 who starts screening in her 40s, two who start in their 50s and three who start in their 60s will avoid a breast cancer death, Dr. Harris said.

Dr. Wender added that while improved treatments clearly helped lower the breast cancer death rate, so did mammography, by catching cancers early.

But an editorial accompanying the new study said that earlier studies that found mammograms helped women were done before the routine use of drugs like tamoxifen that sharply reduced the breast cancer death rate. In addition, many studies did not use the gold-standard methods of the clinical trial, randomly assigning women to be screened or not, noted the editorial's author, Dr. Mette Kalager, and other experts.

Dr. Kalager, an epidemiologist and screening researcher at the University of Oslo and the Harvard School of Public Health, said there was a reason the results were unlike those of earlier studies. With better treatments, like tamoxifen, it was less important to find cancers early. Also, she said, women in the Canadian study were aware of breast cancer and its dangers, unlike women in earlier studies who were more likely to ignore lumps.

"It might be possible that mammography screening would work if you don't have any awareness of the disease," she said.

The Canadian study reached the same conclusion about the lack of a benefit from mammograms after 11 to 16 years of follow-up, but some experts predicted that as time went on the advantages would emerge.

That did not happen, but with more time the researchers could, for the first time, calculate the extent of overdiagnosis, finding cancers that would never have killed the women but that led to treatments that included surgery, chemotherapy and radiation.

Many cancers, researchers now recognize, grow slowly, or not at all, and do

not require treatment. Some cancers even shrink or disappear on their own. But once cancer is detected, it is impossible to know if it is dangerous, so doctors treat them all.

If the researchers also included a precancerous condition called ductal carcinoma in situ, the overdiagnosis rate would be closer to one in three cancers, said Dr. Anthony B. Miller of the University of Toronto, the lead author of the paper. Ductal carcinoma in situ, or D.C.I.S., is found only with mammography, is confined to the milk duct and may or may not break out into the breast. But it is usually treated with surgery, including mastectomy, or removal of the breast.

Mammography's benefits have long been debated, but no nations except Switzerland have suggested the screening be halted. In a recent report, the Swiss Medical Board, an expert panel established by regional ministers of public health, advised that no new mammography programs be started in that country and that those in existence have a limited, though unspecified, duration. Ten of 26 Swiss cantons, or districts, have regular mammography screening programs.

Dr. Peter Juni, a member of the Swiss Medical Board until recently, said one concern was that mammography was not reducing the overall death rate from the disease, but increasing overdiagnosis and leading to false positives and needless biopsies.

"The mammography story is just not such an easy story," said Dr. Juni, a clinical epidemiologist at the University of Bern.

Even experts like Dr. H. Gilbert Welch, a professor of medicine at Dartmouth, who have questioned mammography's benefits were surprised by Switzerland's steps to reconsider its widespread use.

"Wow, times they are a-changin'," Dr. Welch said.

In the United States, about 37 million mammograms are performed annually at a cost of about \$100 per mammogram. Nearly three-quarters of women age 40 and over say they had a mammogram in the past year. More than 90 percent of women ages 50 to 69 in several European countries have had at least one mammogram.

Dr. Kalager, whose editorial accompanying the study was titled "Too Much Mammography," compared mammography to prostate-specific antigen screening for prostate cancer, using data from pooled analyses of clinical trials. It turned out that the two screening tests were almost identical in their overdiagnosis rate and had almost the same slight reduction in breast or prostate

deaths.

“I was very surprised,” Dr. Kalager said. She had assumed that the evidence for mammography must be stronger since most countries support mammography screening and most discourage PSA screening.

A version of this article appears in print on February 12, 2014, on page A1 of the New York edition with the headline: Vast Study Casts Doubts on Value of Mammograms.